JpGU International Symposium 2011 Urgent Session "The 2011 off the Pacific Coast of Tohoku earthquake"

MIS36 P-129

## Inundation area by the 2011 Tohoku earthquake tsunami in Sendai plain: comparison to the 869 Jogan earthquake tsunami



D. Sugawara<sup>1</sup>, F.Imamura<sup>1</sup>, K. Goto<sup>2</sup>, H. Matsumoto<sup>3</sup>, K. Minoura<sup>4</sup>, S. Fujino<sup>5</sup>, T. Haraguchi<sup>6</sup>, K. Fujima<sup>7</sup>, R. Tsudaka<sup>7</sup> and T. Abe<sup>8</sup>



1 Disaster Control Research Center, Graduate School of Engineering, Tohoku University / 2 Planetary Exploration Research Center, Chiba Institute of Technology / 3 Department of Regional Management, Faculty of Liberal Arts, Tohoku Gakuin University / 4 Institute of Geology and Paleontology, Graduate School of Science, Tohoku University / 5 Graduate School of Life and Environmental Sciences, University of Tsukuba / 6 Graduate School of Sciences, Osaka City University / 7 Department of Civil and Environmental Engineering, National Defense Academy of Japan / 8 Department of Geography, Graduate School of Environmental Studies, Nagoya University

## 1. Introduction

The coastal plains in Sendai City and neighboring areas were extensively inundated by the 2011 Tohoku Earthquake Tsunami, and caused massive human and property damages in this region. Data on inundation area in the Sendai Plain (distances and heights of tsunami run-up) have been obtained through the post-tsunami field survey of the 2011 event<sup>[4]</sup>. In this study, inundation area by the 2011 tsunami is compared with that by the Jogan earthquake tsunami, whichi has been known as one of the largest historical event in Tohoku (northeast) region of Japan. Inundation area of the Jogan tsunami was previously estimated based on the sediment distribution and numerical modeling of the tsunami<sup>[1]</sup>.

