# 8. Summary

## Size of the 2011 Tohoku Earthquake Tsunami

Major run-up limits by the 2011 tsunami reached around the numerically-estimated inundation limit of the 869 tsunami, although the coastline has been shifted  $\sim 1$  km seaward.

Inundation area by the 2011 tsunami exceeds that by the 869 tsunami, if minor run-up is taken into acoount.

Judging from the field data in the Sendai Plain, the size of the 2011 tsunami is at least same as or more possibly greater than that of the 869 tsunami.

### **References and data source**

[1] Sugawara, D., Imamura, F., Matsumoto, H., Goto, K. and Minoura, F. (2011) Journal of Japan Society of Natural Disaster Science 29 (4), 501-516 (in Japanese). [2] Geospatial Information Authority of Japan, "Crustal Deformation and Fault Model obtained from GEONET data analysis",

(http://www.gsi.go.jp/cais/topic110422-index-e.html) retrieved on 24 May 2011.

[3] Association of Japanese Geographers, "Maps of the Area hit by the Tsunami of 11 March 2011, Northeast Japan", (http://danso.env.nagoya-u.ac.jp/20110311/map/index e.html) (retrieved on 24 May 2011).

[4] The 2011 Tohoku Earthquake Tsunami Joint Survey Group, "Preliminary results of field survey", (http://www.coastal.jp/tsunami2011/) (retrieved on 24 May 2011).

[5] Watanabe, H. (2001) Historical Earthquakes (Rekishi-jishin) 17, 130–146 (in Japanese).

[6] Haraguchi, T., Torii, K., Fujiwara, O., Shimazaki, K. and Imaizumi, T. (2006) Chikyu 28 (8), 539-545 (in Japanese).

[7] Shishikura, M., Sawai, Y., Okamura, Y., Komatsubara, J., Aung, T.T., Ishiyama, T., Fujiwara, O. and Fujino, S. (2007) Annual Report of Active Fault and Paleoearthquake Researches 7, 31–46 (in Japanese with English Abstract).

[8] Abe, H., Sugeno, Y. and Chigama, A. (1990) Journal of the Seismological Society of Japan (Zishin) 43, 513–525 (in Japanese with English abstract). [9] Minoura, K. and Nakaya, S. (1991) Journal of Geology 99, 265–287.

[10] Minoura, K., Imamura, F., Sugawara, D., Kono, Y. and Iwashita, T. (2001) Journal of Natural Disaster Science 23, 83–88. [11] Sawai, Y., Shishikura, M., Okamura, Y., Takada, K., Matsu' ura, T., Aung, T.T., Komatsubara, J., Fujii, Y., Fujiwara, O., Satake, K., Kamataki, T. and Sato, N. (2007) Annual Report of Active Fault and Paleoearthquake Research 7, 47–80 (in Japanese with English Abstract). [12] Sawai, Y., Shishikura, M. and Komatsubara, J. (2008) Annual Report of Active Fault and Paleoearthquake Researches 8, 17–70 (in Japanese with English Abstract).